

10 Warning Signs of Primary Immunodeficiency

Primary Immunodeficiency (PI) causes children and young adults to have infections that come back frequently or are unusually hard to cure. In America alone, up to 1/2 million people suffer from one of the 100 known Primary Immunodeficiency diseases. If you or someone you know are affected by two or more of the following warning signs, speak to a physician about the possible presence of an underlying Primary Immunodeficiency.


- | | | | |
|---|---|---|----|
| 1 | Eight or more new ear infections within 1 year. | Recurrent, deep skin or organ abscesses. | 6 |
| 2 | Two or more serious sinus infections within 1 year. | Persistent thrush in mouth or elsewhere on skin, after age 1. | 7 |
| 3 | Two or more months on antibiotics with little effect. | Need for intravenous antibiotics to clear infections. | 8 |
| 4 | Two or more pneumonias within 1 year. | Two or more deep-seated infections. | 9 |
| 5 | Failure of an infant to gain weight or grow normally. | A family history of Primary Immunodeficiency. | 10 |

Presented as a public service by:

Jeffrey Modell Foundation 

This publication was supported by grant funds from the United States

Centers for Disease Control and Prevention (CDC) and the generosity of the following organizations:



For information or referrals contact the Jeffrey Modell Foundation: 1-866-INFO-4-PI • www.info4pi.org

These warning signs were developed by the Jeffrey Modell Foundation Medical Advisory Board • © 2003 Jeffrey Modell Foundation